

Seventh Cranial Nerve Palsy

Disclaimer

SEE ALSO: Herpes Zoster Ophthalmicus, Temporal Arteritis (TA)

DESCRIPTION – Seventh (7th) nerve palsy is a dysfunction of the facial nerve (CN7) which supplies the facial muscles, taste to the anterior 2/3 of the tongue, parasympathetic supply to head and neck ganglia. It can be due to either a central or peripheral lesion with partial or complete facial muscle weakness.

HOW TO ASSESS

Red Flags:

- Suspect upper motor neuron (UMN) (e.g. stroke) lesion if ipsilateral sparing of forehead movements
- Associated neurologic signs require further investigation
- Suspect temporal arteritis (TA) if associated symptoms and age over 60
- Beware potentially blinding corneal exposure if poor lid closure.

Aetiology:

Idiopathic:

Bell's palsy – Isolated unilateral lower motor neuron (LMN) lesion. May be related to herpes simplex virus (HSV) activation. Diagnosis of exclusion. See Table 1.

Inflammation of the peripheral nerve:

Ramsay-Hunt syndrome – Acute facial neuropathy from adjacent varicella zoster (HZV) infection with vesicular rash of the ear canal or oropharynx. In the absence of a rash it is termed zoster sine herpette.

Other causes include: otitis media, infectious mastoiditis, parotitis, sarcoidosis, diabetic mononeuropathy, cytomegalovirus, Epstein-Barr virus, HIV seroconversion.

Compression of peripheral nerve:

Trauma (including surgical injury or post-operative swelling), vestibular schwannoma or skull base tumour, facial nerve tumour, cholesteatoma, parotid tumour.

Damage to facial nucleus, fascicle or motor cortex:

Cortical stroke, pontine infarction, demyelinating disease, local infiltration by tumour, TA.

Neuromuscular pathology:

Myasthenia gravis, myotonic dystrophy, Guillain-Barre syndrome.

On History:

- Past history of facial weakness, stroke, malignancy
- Recent trauma to head or face, or recent viral infection
- Facial droop/weakness: evolution. Bell's palsy usually comes on quickly (over hours, up to a few days)
- Painful watery eye, blurred vision
- Facial or ear pain
 - Severe pain is suggestive of HZV
 - Mild pain in the face or behind the ear may occur in Bell's palsy
- Hyperacusis and/or altered taste
 - Suggestive of a lesion proximal to the chorda tympani
 - Both are common in Bell's palsy
- Other neurological symptoms
 - Limb weakness/parasthesia
 - Diplopia – consider intracranial tumours causing palsy of cranial nerves III, IV, VI
 - Deafness, tinnitus, vertigo – consider Ramsay-Hunt, rarely cerebellopontine angle tumour (e.g. vestibular schwannoma)
 - Sensory loss – consider cranial nerve V lesion
- Symptoms of TA: headache, jaw claudication scalp tenderness, polymyalgia rheumatica, weight loss, etc. Refer to TA CPG

On Examination:

- Facial muscle weakness
 - House-Brackmann score (see Table 2)
- Ocular examination
 - Visual acuity, pupils
 - Extra-ocular movements
 - Lids/Cornea
 - Orbicularis function – lid closure
 - Document lagophthalmos on gentle and forced lid closure (mm)
 - Levator function
 - Bell's phenomenon (eyeball rolls upward on attempted eyelid closure by patient in normal Bell's phenomenon)
 - Corneal exposure: conjunctival injection, fluorescein staining-punctate epithelial erosions (PEE), epithelial defect
 - Corneal sensation (V1)
 - Temporal artery: non pulsatile, tender – suggestive of TA
- Ear examination
 - Inspect for vesicular rash – suggestive of Ramsay-Hunt syndrome
 - Canal
 - Tympanic membrane and middle ear
- All cranial nerves: including facial sensation (V1,V2,V3)
- Signs of 7th nerve aberrant regeneration (indicates chronicity) - gustatory sweating, co-contraction of ipsilateral facial muscles (synkinesis) or involuntary tearing on the affected side when eating (crocodile tears)
- Upper/lower limb neurological examination

Table 1: Table comparing features typical of Bell’s palsy and features that may suggest alternative pathology

	Features typical in Bell’s palsy	Features suggestive of alternative pathology
Onset	Rapid – over hours to 3 days	Insidious onset
Pain	Mild facial or post-auricular pain only	Headache or severe facial pain
Associated neurology	Isolated unilateral cranial nerve VII palsy (Note hyperacusis and altered taste commonly occur from chorda tympani involvement)	Bilateral disease, diplopia, paraesthesia, other weaknesses, deafness, tinnitus, vertigo
Systemic features	Nil	Headache, loss of weight, night sweats, fever, fatigue, rash
Recovery	Starts to recover over first 3 weeks	No improvement seen within first 4 weeks

Table 2: House-Brackmann Scale for facial nerve grading system

Grade	Findings
I	Normal symmetrical function
II	Complete eye closure with minimal effort Slight asymmetry of face noticed on maximal effort
III	Complete eye closure Strong but asymmetrical mouth movement Obvious weakness, but not disfiguring
IV	Incomplete eye closure Asymmetry of mouth with maximal effort Severe synkinesis, mass movement or spasm
V	Incomplete eye closure Motion barely perceptible Usually no synkinesis, contracture or spasm present
VI	No movement detectable and loss of tone

On Investigation:

- Swab for viral PCR (including HZV) if rash
- Laboratory investigations for TA, sarcoidosis, neuromuscular etiology if history suggestive
- Consider CT scan if recent history of head or facial trauma, facial mass or surgery
- Consider MRI brain if suspect central lesion. Suggestive clinical features:
 - Gradual onset greater than a few days
 - Sparing of frontalis muscle
 - Any other neurological deficit
- Please remember Bell’s palsy is a diagnosis of exclusion

Management:

Note: all patients must be seen by ophthalmology and ENT

Systemic:

- Steroids – Prednisolone 1 mg/kg orally (up to 75 mg) daily for 5 days
 - Indications:
 - Ramsay-Hunt syndrome
 - Bell’s palsy
 - Other inflammatory pathologies (in consultation with ENT unit)
 - Steroids should be initiated within 48 - 72 hours of onset of symptoms
- Antivirals – Valacyclovir 1,000 mg TDS for 7 days
 - Indications:
 - Ramsay-Hunt syndrome
 - Bell’s palsy (Optional)
 - Note: Routine antiviral use for Bell’s has been proposed due to possible association with herpes simplex infection, however this is controversial.

Ocular:

- Consider for patients with severe facial weakness (House-Brackmann IV-VI)
- All patients with lagophthalmos:
 - Preservative free ocular lubricants (frequency and viscosity based on degree of corneal exposure)
 - Nightly: paraffin based lubricating ointment
 - Consider lid closure with tape/pad. Please note that this can be counterproductive as it is difficult to achieve complete eyelid closure with tape/pad.

Table 3: Corneal exposure risk

Corneal Exposure Risk	Possible Signs	Suggested Management
Mild	Few PEE Eyes white Complete forced lid closure Minimal lagophthalmos	Regular ocular lubricant drops (6x per day) +/- Nocte lubricant ointment Avoid environments that aggravate dryness
Moderate	Significant PEE Ocular injection Lagophthalmos 1 – 2 mm	Preservative-free ocular lubricant drops 2-hourly Nocte lubricant ointment Tape lid closed overnight Consider goggles to keep ocular humidity
Severe	Epithelial defect Widespread PEE Lagophthalmos \geq 3 mm Reduced corneal sensation Corneal thinning Poor Bell’s phenomenon	Preservative-free ocular lubricant drops 1-hourly Nocte lubricant ointment Tape lid closed overnight Consider tarsorrhaphy

Dental:

Indications: any patients with inability to close mouth.

Significant oral dryness can result in increased risk of caries.

Prognosis in Bell's palsy:

- Weakness may worsen in the first three weeks
- Most (85%) will start to recover within three weeks. 71% of patients with Bell's palsy have full recovery by 12 months, many of which occur within 3 months
Patients with incomplete paralysis are highly likely to completely recover (94%).
Patients with more severe initial weakness (House-Brackmann III – VI) are likely to have ongoing long-term dysfunction
- Recurrence of Bell's palsy occurs in approximately 7% of patients within 10 years
- Some patients may recover with aberrant facial nerve regeneration

Follow Up:

- Consider referral to Neurology:
 - Atypical features
 - No improvement is observed within first 4 weeks
 - Incomplete resolution of symptoms by 3 months
- Review Acute ENT in 4 weeks (for House-Brackmann III-VI)
- Ophthalmology review (AOS) in 1-2 weeks depending on the degree of lagophthalmos and corneal exposure

Authors: Dr Lukas Sahhar, Dr Kristen Wells and CPG Working Party

Review Date: 30/05/2023

Evidence Table

Author/s	Title	Source	Level of Evidence (I – VII)	Comments
	Bell's Palsy	UpToDate 2017	I	
	Bell's Palsy 2013	American Academy of Otolaryngology Head and Neck Surgery Clinical Practice Guideline	I	
Pane A, Burdon M, Miller N. Mosby	The Neuro-Ophthalmology Survival Guide	Elsevier	II	
	Electronic Therapeutic Guidelines, Bell's Palsy Therapeutic Guidelines Australia		I	

The Hierarchy of Evidence

The Hierarchy of evidence is based on summaries from the National Health and Medical Research Council (2009), the Oxford Centre for Evidence-based Medicine Levels of Evidence (2011) and Melynck and Fineout-Overholt (2011).

I Evidence obtained from a systematic review of all relevant randomised control trials.

II Evidence obtained from at least one well designed randomised control trial.

III Evidence obtained from well-designed controlled trials without randomisation.

IV Evidence obtained from well designed cohort studies, case control studies, interrupted time series with a control group, historically controlled studies, interrupted time series without a control group or with case series.

V Evidence obtained from systematic reviews of descriptive and qualitative studies.

VI Evidence obtained from single descriptive and qualitative studies.

VII Expert opinion from clinician, authorities and/or reports of expert committees or based on physiology.

CPG Suite General Disclaimer

These CPGs were written for use in the RVEEH speciality Emergency Department. They should be used under the guidance of an ENT or Ophthalmology registrar, and certain medications / procedures should only be undertaken by speciality registrars.

If you require clinical advice, please contact our admitting officer for assistance:

EYE: 03 9929 8033 ENT: 03 9929 8032